

A CHARACTERIZATION OF OPERATORS WITH p -SUMMING ADJOINT VIA p -LIMITED SETS

JUAN MANUEL DELGADO AND CÁNDIDO PIÑEIRO

A subset A of a Banach space X is limited if and only if for every weak* null sequence (x_n^*) in X^* there exists a sequence $(\alpha_n) \in c_0$ such that $|\langle x_n^*, x \rangle| \leq \alpha_n$ for all $x \in A$ and $n \in \mathbb{N}$. As an extension of this notion, a subset A of a Banach space X is said to be p -limited ($p \in [1, \infty)$) if for every weakly p -summable sequence (x_n^*) in X^* there exists $(\alpha_n) \in \ell_p$ such that $|\langle x_n^*, x \rangle| \leq \alpha_n$ for all $x \in A$ and $n \in \mathbb{N}$ [2]. Some basic properties related to this notion are showed as well as its connections with (different forms of) compactness. As an application, we give a characterization of operators with p -summing adjoint as those mapping relatively compact sets to p -limited sets.

The research was supported by MTM2009-14483-C02-01 and MTM2012-36740-C02-01 projects (Spain).

REFERENCES

- [1] J. M. DELGADO AND C. PIÑEIRO, *A note on p -limited sets*. preprint.
- [2] A. K. KARN AND D. P. SINHA, *An operator summability in sequences in Banach spaces*, arXiv:1207.3620 [math.FA] (2012).

HIGHER TECHNICAL SCHOOL OF ARCHITECTURE, UNIVERSITY OF SEVILLE,
41012 SEVILLE, AV. REINA MERCEDES, 2, SPAIN
E-mail address: `jmdelga@us.es`

FACULTY OF EXPERIMENTAL SCIENCES, UNIVERSITY OF HUELVA, 21071 HUELVA,
CAMPUS UNIVERSITARIO DE EL CARMEN, SPAIN
E-mail address: `candido@uhu.es`